arXiv:1305.0474v1 [gr-qc] 2 May 2013

Scaling attractors in interacting teleparallel dark energy

G. Otalora

Instituto de Física Teórica, UNESP-Univ Estadual Paulista
Caixa Postal 70532-2, 01156-970 São Paulo, Brazil

Abstract. It has been proposed recently the existence of a non-minimal coupling between a canonical scalar field (quintessence) and gravity in the framework of teleparallel gravity, motivated by similar constructions in the context of General Relativity. The dynamics of the model, known as teleparallel dark energy, has been further developed, but no scaling attractor has been found. Here we consider a model in which the non-minimal coupling is ruled by a dynamically changing $\alpha \equiv f_{,\phi}/\sqrt{f}$, with $f(\phi)$ an arbitrary function of the scalar field ϕ. It is shown that in this case the existence of scaling attractors is possible, which means that the universe will eventually enter these scaling attractors, regardless of the initial conditions. As a consequence, the cosmological coincidence problem could be alleviated without fine-tunings.

1 Introduction

Since the discovery of the universe accelerated expansion, this issue has been one of the most active fields in modern cosmology. The simplest explanation for the dark energy—the name given to the agent responsible for such accelerated expansion—is provided by a cosmological constant. However, this scenario is plagued by a severe fine tuning problem associated with its energy scale [1, 2, 3]. This problem can be alleviated by considering a scalar field with a dynamically varying equation of state, like for example quintessence dark energy models. Scalar fields naturally arise in particle physics (including string theory), and these fields can act as candidates for dark energy. So far a wide variety of scalar-field dark energy models have been proposed, such as quintessence, phantoms, k-essence, amongst many others [1, 4].

Usually, dark energy models are based on scalar fields minimally coupled to gravity, as well as a possible coupling of the field to a background fluid (dark matter). Also, a possible coupling between the scalar field and the Ricci scalar R is not to be excluded in the context of generalized Einstein gravity theories. For example, a quintessence field coupled to gravity and called "extended quintessence" was proposed in Ref. [5] (see also Refs. [6, 7]). K-essence models non-minimally coupled to gravity were studied in Refs. [8, 9].

On the other hand, the Teleparallel Equivalent of General Relativity (TEGR) [10, 11, 12] is an equivalent formulation of classical gravity, in which instead of using the torsionless Levi-Civita connection one uses the curvatureless Weitzenböck one. The dynamical objects are the four linearly independent tetrads, not the metric. The advantage of this framework is that the torsion tensor is formed solely from products of first derivatives of the tetrad. Thus, apart from some conceptual differences, TEGR is completely equivalent and indistinguishable from General Relativity (GR) at the level of field equations [11, 12]. Recently the teleparallel gravity generalization, the so-called $F(T)$ theory [13, 14], has attracted much attention.

In Refs. [15, 16] it was proposed a non-minimal coupling between quintessence and gravity in the framework of teleparallel gravity, motivated by a similar construction in the context of GR. The theory was called "teleparallel dark energy", and its dynamics was studied later in [17, 18]. However, no scaling attractor was found. Here we generalize the non-minimal coupling function $\phi^2 \to f(\phi)$ and we define $\alpha \equiv f_{,\phi}/\sqrt{f}$ such that α is constant for $f(\phi) \propto \phi^2$, but it can also be a dynamically changing quantity $\alpha(\phi)$. For dynamically changing α we show the existence of scaling attractors with accelerated expansion in agreement with observations. As long as the scaling solution is an attractor, for any generic initial conditions the system would sooner or later enter the scaling regime in which the dark energy density is comparable to the matter energy density with accelerated expansion of the universe, alleviating in this the fine tuning problem of dark energy [1, 4].

2 Quintessence field in general relativity

As is well known [5, 6, 7], one can generalize quintessence by including a non-minimal coupling between quintessence and gravity. The relevant action reads

$$S = \int d^4x \sqrt{-g}\left[\frac{R}{2\,\kappa^2} + \frac{1}{2}\,\partial_\mu\phi\,\partial^\mu\phi - V(\phi) + \xi\ f(\phi)\,R\right] + S_m, \tag{1}$$

where R is the Ricci scalar, S_m is the matter action, $\kappa^2 = 8\pi G$ and $c = 1$ (we use the metric signature convention $(+,-,-,-)$ throughout). Also, the parameter ξ is a dimensionless constant and $f(\phi)$ (with units of $mass^2$ and positive) is an arbitrary function of ϕ responsible for non-minimal coupling. In [19, 20, 21, 22] adopted $f \backsim \phi^2$ in the context inflation, other forms have been considered in [23, 24] (for a more general form that includes derivatives see [25]). When including a non-minimal coupling in quintessence, the corresponding Friedmann equations are preserved. On the other hand, the pressure density and energy density of quintessence are changed to

$$p_\phi = \frac{1}{2}\left(1 + 4\,\xi\,f_{,\phi\phi}\right)\dot{\phi}^2 - V + 4\,\xi\left(f + 3\,\xi\,f_{,\phi}^2\right)\dot{H} - \\ 2\,\xi\,f_{,\phi}\,\dot{\phi}\,H - 2\,\xi\,f_{,\phi}\,V_{,\phi} + 6\,\xi\left(f + 4\,\xi\,f_{,\phi}^2\right)H^2, \tag{2}$$

$$\rho_\phi = \frac{1}{2}\,\dot{\phi}^2 + V - 6\,\xi\,H\,f_{,\phi}\,\dot{\phi} - 6\,\xi\,H^2\,f(\phi), \tag{3}$$

where $V_{,\phi} \equiv dV/d\phi$ and $f_{,\phi} \equiv df/d\phi$. For find p_ϕ and ρ_ϕ we have used the equation of motion

$$\ddot{\phi} + 3\,H\,\dot{\phi} - \xi\,R\,f_{,\phi} + V_{,\phi} = 0, \tag{4}$$

calculated from the action (1), and the useful expression $R = 6\left(\dot{H} + 2\,H^2\right)$ in the flat Friedmann-Robertson-Walker (FRW) geometry, see [6, 7].

3 Teleparallel equivalent to general relativity

Following [12], we briefly review the key ingredients of teleparallel gravity. The orthonormal tetrad components $h_a(x^\mu)$ relate to the metric through

$$g_{\mu\nu} = h^a{}_\mu\,h^b{}_\nu\,\eta_{ab} \tag{5}$$

where Latin indices $a, b, c, \ldots$ are indices running over $0, 1, 2, 3$ for the tangent space of the manifold, and Greek indices $\mu, \nu, \rho, \ldots$ are the coordinate indices on the manifold, also running over $0, 1, 2, 3$. In teleparallel gravity, the relevant action is

$$S = \frac{1}{2\,\kappa^2} \int dx^4\, h\, T + S_m \tag{6}$$

where $h = \det(h^a{}_\mu) = \sqrt{-g}$. The torsion scalar T is given by

$$T \equiv S_\rho{}^{\mu\nu}\, T^\rho{}_{\mu\nu}, \tag{7}$$

where

$$T^\rho{}_{\mu\nu} \equiv h_a{}^\rho \left(\partial_\mu h^a{}_\nu - \partial_\nu h^a{}_\mu\right), \tag{8}$$

is the torsion tensor

$$K^{\mu\nu}{}_\rho \equiv \left(T^{\mu\nu}{}_\rho - T^{\nu\mu}{}_\rho - T_\rho{}^{\mu\nu}\right), \tag{9}$$

is the contortion tensor and

$$S_\rho{}^{\mu\nu} \equiv \frac{1}{2}\left(K^{\mu\nu}{}_\rho + \delta^\mu_\rho\, T^{\theta\nu}{}_\theta - \delta^\nu_\rho\, T^{\theta\mu}{}_\theta\right), \tag{10}$$

is the superpotential. Variation with respect to the tetrad fields yields the teleparallel version of the gravitational field equation

$$h^{-1}\,\partial_\mu\left(h\, h_a{}^\rho\, S_\rho{}^{\mu\nu}\right) - h_a{}^\lambda\, T^\rho{}_{\mu\lambda}\, S_\rho{}^{\nu\mu} - \frac{1}{4}\, h_a{}^\nu\, T = \frac{\kappa^2}{2}\, h_a{}^\rho\, \Theta_\rho{}^\nu \tag{11}$$

where $\Theta_\rho{}^\nu$ stands for the usual energy-momentum tensor. These equations are exactly the same as those of GR for every geometry choice. In the case of a flat FRW background metric, the tetrad choice have the form

$$h^a{}_\mu(t) = \mathrm{diag}(1, a(t), a(t), a(t)), \tag{12}$$

where $a(t)$ is the scale factor. It is an exact solution of the field equation (11) and it is seen that the corresponding Friedmann equations are identical to the GR ones both at the background and perturbation levels [1, 2, 11, 15].

4 Interacting teleparallel dark energy

In what follows we consider a non-minimal coupling between quintessence field and teleparallel gravity. The action will simply read

$$S = \int d^4x\, h \left[\frac{T}{2\,\kappa^2} + \frac{1}{2}\,\partial_\mu\phi\,\partial^\mu\phi - V(\phi) + \xi\, f(\phi)\, T\right] + S_m. \tag{13}$$

Also, we consider a possible interaction between teleparallel dark energy and dark matter as was studied in [17], since there is no physical argument to exclude the possible interaction between them. Varying the action (13) with respect to the tetrad fields yields

$$\begin{aligned} 2\left(\frac{1}{\kappa^2} + 2\,\xi\, f(\phi)\right)&\left[h^{-1}\, h^a{}_\alpha\, \partial_\sigma\left(h\, h_a{}^\tau\, S_\tau{}^{\rho\sigma}\right) + T^\tau{}_{\nu\alpha}\, S_\tau{}^{\rho\nu} + \frac{T}{4}\,\delta_\alpha{}^\rho\right] \\ &+ 4\,\xi\, S_\alpha{}^{\rho\sigma}\, f_{,\phi}\,\partial_\sigma\phi + \left[\frac{1}{2}\,\partial_\mu\phi\,\partial^\mu\phi - V(\phi)\right]\delta_\alpha{}^\rho - \partial_\alpha\phi\,\partial^\rho\phi = \Theta_\alpha{}^\rho. \end{aligned} \tag{14}$$

Therefore, imposing the flat FRW geometry (12) in (14), we obtain the Friedmann equations with

$$\rho_\phi = \frac{1}{2}\,\dot\phi^2 + V(\phi) - 6\,\xi\,H^2\,f(\phi), \tag{15}$$

the scalar energy density and

$$p_\phi = \frac{1}{2}\,\dot\phi^2 - V(\phi) + 4\,\xi\,H\,f_{,\phi}\,\dot\phi + 6\,\xi\,\left(3\,H^2 + 2\,\dot H\right)\,f(\phi), \tag{16}$$

the pressure density of field. Here we also use the useful relation $T = -6\,H^{\,2}$, which arises for flat FRW geometry.

In the flat FRW background, the variation of the action (13) with respect to scalar field yields the field equation

$$\ddot\phi + 3\,H\,\dot\phi + V_{,\phi} + 6\,\xi\,f_{,\phi}\,H^2 = -\sigma. \tag{17}$$

where σ is the scalar charge corresponds to coupling between dark matter and dark energy, and it is defined by the relation $\delta S_m/\delta\phi = -h\,\sigma$ [1, 4, 30]. Rewriting (17) in terms of ρ_ϕ and p_ϕ we find the continuity equation for the field

$$\dot\rho_\phi + 3\,H\,\rho_\phi\,(1+\omega_\phi) = -Q, \tag{18}$$

whereas for matter

$$\dot\rho_m + 3\,H\,\rho_m\,(1+\omega_m) = Q, \tag{19}$$

where $\omega_\phi \equiv p_\phi/\rho_\phi$ and $\omega_m \equiv p_m/\rho_m = const \geq 0$ are the equation-of-state parameter of dark energy and dark matter, respectively. We define the barotropic index $\gamma \equiv 1 + \omega_m$ such that $1 \leq \gamma < 2$. The parameter $Q \equiv \dot\phi\,\sigma$ is the coupling between dark energy and dark matter.

5 Dynamical system of interacting teleparallel dark energy

Of particular importance in the investigation of cosmological scenarios are those solutions in which the energy density of the scalar field mimics the background fluid energy density, $\rho_\phi/\rho_m = C$, with C a nonzero constant. Cosmological solutions which satisfy this condition are called "scaling solutions" and the cosmological coincidence problem can be alleviated in most dark energy models via scaling attractors (it is defined later) [1, 4]. To study the cosmological dynamics of the model, we introduce the followings dimensionless variables:

$$x \equiv \frac{\kappa\,\dot\phi}{\sqrt{6}\,H} \qquad y \equiv \frac{\kappa\,\sqrt{V}}{\sqrt{3}\,H}, \qquad u \equiv \kappa\,\sqrt{f(\phi)}, \qquad \lambda \equiv -\frac{V_{,\phi}}{\kappa\,V}, \qquad \alpha \equiv \frac{f_{,\phi}}{\sqrt{f}}. \tag{20}$$

Using these variables we define

$$s \equiv \frac{\dot H}{H^2} = \frac{-3\,\gamma\,y^2 + 3\,(2-\gamma)\;x^2 + 4\,\sqrt{6}\,\alpha\,\xi\,u\,x}{2\;(2\,\xi\,u^2+1)} + \frac{3\,\gamma}{2}, \tag{21}$$

and the Eqs. (18) and (19) can be rewritten as a dynamical system of ordinary differential equations (ODE), namely

$$x' = \frac{\sqrt{6}}{2}\,\lambda\,y^2 - \sqrt{6}\,\alpha\,\xi\,u + x\;(s-3) - \hat{Q}, \tag{22}$$

$$y' = \left(s - \frac{\sqrt{6}\,\lambda\,x}{2} \right) y, \tag{23}$$

$$u' = \frac{\sqrt{6}\,\alpha\,x}{2}, \tag{24}$$

$$\lambda' = -\sqrt{6}\,\left(\Gamma - 1\right)\,\lambda^2\,x, \tag{25}$$

$$\alpha' = \sqrt{6}\,\left(\Pi - \frac{1}{2}\right)\,\frac{\alpha^2\,x}{u}. \tag{26}$$

In these equations primes denote derivative with respect to the so-called e-folding time $N \equiv \ln a$. Also, we define

$$\hat{Q} \equiv \frac{\kappa\,Q}{\sqrt{6}\,H^2\,\dot{\phi}}, \qquad \Pi \equiv \frac{f\,f_{,\phi\phi}}{f_{,\phi}^2}, \qquad \Gamma \equiv \frac{V\,V_{,\phi\phi}}{V_{,\phi}^2}. \tag{27}$$

The fractional energy densities $\Omega \equiv (\kappa^2 \rho)/(3H^2)$ for the scalar field and background matter are given by

$$\Omega_\phi = y^2 + x^2 - 2\,\xi\,u^2, \qquad \Omega_m = 1 - \Omega_\phi, \tag{28}$$

respectively. The equation of state of the field $\omega_\phi = p_\phi/\rho_\phi$ reads

$$\omega_\phi = \frac{x^2 - y^2 + 4\,\sqrt{\frac{2}{3}}\,\alpha\,\xi\,u\,x + 2\left(1 - \frac{2}{3}\,s\right)\,\xi\,u^2}{(y^2 + x^2) - 2\,\xi\,u^2}. \tag{29}$$

On the other hand, the effective equation of state $\omega_{eff} = \left(p_m + p_\phi\right)/\left(\rho_m + \rho_\phi\right)$ is given by

$$\omega_{eff} = -\,\gamma\,y^2 - (\gamma - 2)\,x^2 + 4\,\sqrt{\frac{2}{3}}\,\alpha\,\xi\,u\,x + 2\,\xi\left(\gamma - \frac{2}{3}\,s\right)\,u^2 + \gamma - 1, \tag{30}$$

and the accelerated expansion occurs for $\omega_{eff} < -1/3$.

Here, we should emphasize that the dynamical system of ordinary differential equations (ODE) equations (22)-(26) are not a dynamical autonomous system unless the parameters Γ and Π are know. From now on we concentrate on exponential scalar-field potential of the form $V(\phi) = V_0\,e^{-\lambda\,\kappa\,\phi}$, such that λ is a positive constant, that is, $\Gamma = 1$ (equivalently, we could consider potentials satisfying $\lambda \equiv -V_{,\phi}/\kappa\,V \approx const$, which is valid for arbitrary but nearly flat potentials [26, 27, 28]). In addition to the fact that exponential potentials can give rise to an accelerated expansion (a power-law expansion $a(t) \propto t^p$ with $p > 1$), they possess cosmological scaling solutions [1, 4]. On the other hand, for a general coupling function $u \equiv \kappa\,\sqrt{f(\phi)}$, with inverse function $\phi = f^{-1}(u^2/\kappa^2)$, $\alpha(\phi)$ and $\Pi(\phi)$ can be expressed in terms of u (this approach is similar to that followed in the case of quintessence in GR with potential beyond exponential potential, see [29]). In this case, when α and Π are functions of u, the four-dimensional autonomous system (22)-(26) ($\lambda = const$) reduces to three-dimensional autonomous system (22)-(24) with $\alpha = \alpha(u)$. Combining the equations (24) and (26) we find that

$$\alpha(u) = C\,e^{\int \frac{2\,\Pi(u) - 1}{u}\,d\,u}, \tag{31}$$

where C is nonzero constant. For example for $\Pi = 1/2$ then $\alpha = const \neq 0$. On the other hand, if for instance $(2\,\Pi(u) - 1)\,/u \propto 1/\,(u - u_c)$ then $\alpha(u) \propto\ (u - u_c) \to 0$ and $\alpha^2(u)\,\Pi(u) \to 0$ when $u \to u_c$ with u_c a dimensionless constant ($\alpha(u) \to const = 0$). Also, in this case $f(\phi)$ is calculated from (20) and $f(\phi) \to u_c^2/\kappa^2 = const$ when $\phi \to \pm\infty$. Therefore, in general, two situations may arise; one where α is a nonzero constant and another where α depends on u with an asymptotic behavior $\alpha(u) \to \alpha(u_c) = 0$ when $u \to u_c$ and u_c a dimensionless constant. In any case we have a three-dimensional autonomous system (22)-(24) (with $\lambda = const \geq 0$). Once identified the autonomous system we can obtain the fixed points or critical points (x_c, y_c, u_c) by imposing the conditions $x_c' = y_c' = u_c' = 0$. From the definition (20), x_c, y_c, u_c should be real, with $y_c \geq 0$ and $u_c \geq 0$. To study the stability of the critical points, we substitute linear perturbations, $x \to x_c + \delta x$, $y \to y_c + \delta y$, and $u \to u_c + \delta u$ about the critical point (x_c, y_c, u_c) into the autonomous system (22)-(24) and linearize them. The eigenvalues of the perturbations matrix $\mathcal{M}$, namely, μ_1, μ_2 and μ_3, determine the conditions of stability of the critical points. One generally uses the following classification [1, 4]: (i) Stable node: $\mu_1 < 0$, $\mu_2 < 0$ and $\mu_3 < 0$. (ii) Unstable node: $\mu_1 > 0$, $\mu_2 > 0$ and $\mu_3 > 0$. (iii) Saddle point: one or two of the three eigenvalues are positive and the other negative. (iv) Stable spiral: The determinant of the matrix $\mathcal{M}$ is negative and the real parts of μ_1, μ_2 and μ_3 are negative. A critical point is an attractor in the cases (i) and (iv), but it is not so in the cases (ii) and (iii). The universe will eventually enter these attractor solutions regardless of the initial conditions. Now we are going to study the autonomous dynamical system (22)-(24), first for $\alpha = const \neq 0$ and then for dynamically changing $\alpha(u)$, such that $\alpha(u) \to \alpha(u_c) = 0$ when the system falls into the critical point (x_c, y_c, u_c). Also, the fact that the energy density of dark energy (Ω_ϕ) is the same order as that of dark matter (Ω_m) in the present universe, suggests that there may be some relation or interaction between them. Several different forms of the coupling between dark energy and dark matter have been proposed. One possibility usually studied is to consider an interaction of the form $Q = \beta\,\kappa\,\rho_m\,\dot{\phi}$ [1, 4, 30]. A second approach is to introduce an interaction of the form $\Upsilon\,\rho_m$ on the r.h.s. of the continuity equations with the normalization of Υ in terms of the Hubble parameter H, i.e. $\Upsilon/H = \tau$, where τ is a dimensionless coupling [4]. Here we consider only the first possibility ($Q = \beta\,\kappa\,\rho_m\,\dot{\phi}$), for both, constant α and dynamically changing α. In what follows we shall restrict our analysis to the case of $\beta \geq 0$ ($\beta = 0$ case without coupling), but it is straightforward to extend the analysis to the case of negative β.

6 Constant α

6.1 Critical points

From Eq. (27), for the coupling $Q = \beta\,\kappa\,\rho_m\,\dot{\phi}$ it is easy to find that

$$\hat{Q} = \frac{\sqrt{6}\,\beta\,\Omega_m}{2}, \tag{32}$$

with Ω_m given in the equation (28). Substituting (32) in (22) we obtain the critical points of the autonomous system (22)-(24) for constant $\alpha \neq 0$. The critical points are presented in the table 1, and these are a generalization of the fixed points that were found in [17, 18]. In table 2 we summarize the stability properties (to be studied below), and conditions for acceleration and existence for each point. The parameters $q_\pm$ are defined as

$$q_\pm = -\alpha\,\xi \pm \sqrt{\xi\ (\alpha^2\,\xi - 2\,\beta^2)}, \tag{33}$$

Table 1: Critical Points for the autonomous system (22)-(24) for constant $\alpha \neq 0$.

Name	x_c	y_c	u_c	Ω_ϕ	ω_ϕ	ω_{eff}
I.a	0	0	0	0	-1	$\gamma-1$
I.b	0	0	$\frac{q_-}{2\,\beta\,\xi}$	$-\frac{q_-^2}{2\,\beta^2\,\xi}$	$\gamma-1$	$\gamma-1$
I.c	0	0	$\frac{q_+}{2\,\beta\,\xi}$	$-\frac{q_+^2}{2\,\beta^2\,\xi}$	$\gamma-1$	$\gamma-1$
I.d	0	$\frac{\sqrt{\alpha\,v_-}}{\lambda}$	$\frac{v_-}{2\,\lambda\,\xi}$	1	-1	-1
I.e	0	$\frac{\sqrt{\alpha\,v_+}}{\lambda}$	$\frac{v_+}{2\,\lambda\,\xi}$	1	-1	-1

Table 2: Stability properties, and conditions for acceleration and existence of the fixed points in table 1.

Name	Stability	Acceleration	Existence
I.a	Saddle	No	$\beta = 0$
I.b	Saddle	No	$\xi < 0$, $\alpha > 0$ and $\beta > 0$
I.c	Saddle	No	No
I.d	Stable node or spiral, or saddle	All values	$\xi \geq \frac{2\,\lambda^2}{\alpha^2}$, $\alpha > 0$ or $\xi < 0$, $\alpha < 0$
I.e	Saddle	All values	$\xi \geq \frac{2\,\lambda^2}{\alpha^2}$ and $\alpha > 0$

and the parameters $v_\pm$ as

$$v_\pm = \alpha\,\xi \pm \sqrt{\xi\,\left(\alpha^2\,\xi - 2\,\lambda^2\right)}. \tag{34}$$

The point I.a is a matter-dominated solution ($\Omega_m = 1$) with equation of state type cosmological constant $\omega_\phi = -1$, that exists for all values of ξ, λ and $\beta = 0$. Point I.b is a scaling solution and exists solely for $\xi < 0$, $\alpha > 0$ and $\beta > 0$, since in this case the condition $0 < \Omega_\phi < 1$ is satisfied. Accelerated expansion never occurs for these two points because $\omega_{eff} > -1/3$. The critical point I.c is define for $\xi \geq 2\,\beta^2/\alpha^2$, $\alpha < 0$ and $\beta > 0$, but, it is not a physically meaningful solution since $\Omega_\phi < 0$. The points I.d and I.e both correspond to dark-energy-dominated de Sitter solutions with $\Omega_\phi = 1$ and $\omega_\phi = \omega_{eff} = -1$. Point I.d exist for $\xi \geq 2\,\lambda^2/\alpha^2$ and $\alpha > 0$ or $\xi < 0$ and $\alpha < 0$. Point I.e exist for $\xi \geq 2\,\lambda^2/\alpha^2$ and $\alpha > 0$. We note that the points I.d and I.e exist irrespective of the presence of the coupling β, since $\Omega_\phi = 1$ in these cases.

6.2 Stability

Substituting the linear perturbations, $x \to x_c + \delta x$, $y \to y_c + \delta y$, and $u \to u_c + \delta u$ into the autonomous system (22)-(24) and linearize them, we find the follows components for the matrix of linear perturbations $\mathcal{M}$

$$\mathcal{M}_{11} = \frac{-3\,\gamma\,y_c^2 + 9\,(2-\gamma)\,x_c^2 + 8\,\sqrt{6}\,\alpha\,\xi\,u_c\,x_c}{2\,(2\,\xi\,u_c^2 + 1)} - \frac{3\,(2-\gamma)}{2} - \frac{\partial\hat{Q}}{\partial x}|_{x_c,y_c,u_c}, \tag{35}$$

$$\mathcal{M}_{12} = \sqrt{6}\,\lambda\,y_c - \frac{3\,\gamma\,x_c\,y_c}{2\,\xi\,u_c^2 + 1} - \frac{\partial\hat{Q}}{\partial y}|_{x_c,y_c,u_c}, \tag{36}$$

$$\mathcal{M}_{13} = \frac{6\,u_c\,x_c\,\xi\,\left(\gamma\,y_c^2-(2-\gamma)\,x_c^2\right)+4\,\sqrt{6}\,\alpha\,\xi\,x_c^2}{\left(2\,\xi\,u_c^2+1\right)^2} - \frac{2\,\sqrt{6}\,\alpha\,\xi\,x_c^2}{2\,\xi\,u_c^2+1} - \sqrt{6}\,\alpha\,\xi - \frac{\partial\hat{Q}}{\partial u}|_{x_c,y_c,u_c}, \tag{37}$$

$$\mathcal{M}_{21} = \left(\frac{3\,(2-\gamma)\,x_c+2\,\sqrt{6}\,\alpha\,\xi\,u_c}{2\,\xi\,u_c^2+1} - \frac{\sqrt{6}\,\lambda}{2}\right) y_c, \tag{38}$$

$$\mathcal{M}_{22} = \frac{-9\,\gamma\,y_c^2+3\,(2-\gamma)\,x_c^2+4\,\sqrt{6}\,\alpha\,\xi\,u_c\,x_c}{2\,\left(2\,\xi\,u_c^2+1\right)} - \frac{\sqrt{6}\,\lambda\,x_c-3\,\gamma}{2}, \tag{39}$$

$$\mathcal{M}_{23} = \frac{6\,u_c\,y_c\,\xi\,\left(\gamma\,y_c^2+(2-\gamma)\,x_c^2\right)+4\,\sqrt{6}\,\alpha\,\xi\,x_c\,y_c}{\left(2\,\xi\,u_c^2+1\right)^2} - \frac{2\,\sqrt{6}\,\alpha\,\xi\,x_c\,y_c}{2\,\xi\,u_c^2+1}, \tag{40}$$

$$\mathcal{M}_{31} = \frac{\sqrt{6}\,\alpha}{2}, \qquad \mathcal{M}_{32} = 0, \qquad \mathcal{M}_{33} = 0. \tag{41}$$

From the equation (32) we calculate $\partial\hat{Q}/\partial x^i$ and replace in (35)-(37). The eigenvalues of the matrix $\mathcal{M}$ are as follows

- Point I.a:

$$\mu_{1,2} = \frac{3\,(2-\gamma)\,\left(-1\pm\sqrt{1-\frac{16\,\alpha^2\,\xi}{3\,(2-\gamma)^2}}\right)}{4}, \qquad \mu_3 = \frac{3\,\gamma}{2}. \tag{42}$$

- Point I.b:

$$\mu_{1,2} = \frac{3\,(2-\gamma)\,\left(-1\pm\sqrt{1+\frac{16\,\alpha\,\sqrt{\xi\,(\alpha^2\,\xi-2\,\beta^2)}}{3\,(\gamma-2)^2}}\right)}{4}, \qquad \mu_3 = \frac{3\,\gamma}{2}. \tag{43}$$

- Point I.c:

$$\mu_{1,2} = \frac{3\,(2-\gamma)\,\left(-1\pm\sqrt{1-\frac{16\,\alpha\,\sqrt{\xi\,(\alpha^2\,\xi-2\,\beta^2)}}{3\,(\gamma-2)^2}}\right)}{4}, \qquad \mu_3 = \frac{3\,\gamma}{2}. \tag{44}$$

- Point I.d:

$$\mu_{1,2} = \frac{3\,\left(-1\pm\sqrt{1-\frac{4}{3}\,\alpha\,\sqrt{\xi\,\left(\alpha^2\,\xi-2\,\lambda^2\right)}}\right)}{2}, \qquad \mu_3 = -3\,\gamma. \tag{45}$$

- Point I.e:

$$\mu_{1,2} = \frac{3\,\left(-1 \pm \sqrt{1 + \frac{4}{3}\,\alpha\,\sqrt{\xi\,\left(\alpha^2\,\xi - 2\,\lambda^2\right)}}\right)}{2}, \qquad \mu_3 = -3\,\gamma. \tag{46}$$

The Points I.a, I.b, and I.c are saddle points in any case, since $\mu_2 < 0$ and $\mu_3 > 0$. For $\xi < 0$ and $\alpha < 0$ point I.d is a saddle point, but, for

$$\frac{2\,\lambda^2}{\alpha^2} < \xi \leq \frac{\lambda^2}{\alpha^2}\,\left(1 + \sqrt{1 + \frac{9}{16\,\lambda^4}}\right), \tag{47}$$

and $\alpha > 0$ it is a stable node. Also, when $\xi > \frac{\lambda^2}{\alpha^2}\,\left(1 + \sqrt{1 + \frac{9}{16\,\lambda^4}}\right)$ then μ_1 and μ_2 are complex with real part negative and $\det(\mathcal{M}) = -9\,\alpha\,\gamma\,\sqrt{\xi\,\left(\alpha^2\,\xi - 2\,\lambda^2\right)} < 0$ for $\alpha > 0$. Therefore, in this case point I.d is a stable spiral. Finally, point I.e is a saddle point, since it exists solely for $\xi \geq 2\,\lambda^2/\alpha^2$ and $\alpha > 0$.

So, for constant $\alpha > 0$, ξ as in (47) and $\beta = 0$, the universe enters the matter-dominated solution I.a with $\omega_\phi = -1$ and $\omega_{eff} = 0$ for $\gamma = 1$ (non-relativistic dark matter), but since this solution is unstable, finally the universe fall into the attractor I.d (dark-energy-dominated de Sitter solution) with $\Omega_\phi = 1$ and $\omega_\phi = \omega_{eff} = -1$. On the other hand, for $\beta > 0$, $\xi < 0$ and $\alpha > 0$ the system enters in the unstable scaling solution I.b, but finally the solution I.d is not reached since it is a attractor only when $\alpha > 0$ and $\xi > 0$ as in (47). Also, for constant α no scaling attractor was found, even when we allow the possible interaction between teleparallel dark energy and dark matter.

7 Dynamically changing α

Now let us consider a general function of non-minimal coupling $f(\phi)$ such that α depends on u and $\alpha(u) \to \alpha(u_c) = 0$ when $(x, y, u) \to (x_c, y_c, u_c)$ ((x_c, y_c, u_c) is a fixed point of the system). The field ϕ rolls down toward $\pm\infty$ ($x > 0$ or $x < 0$) with $f(\phi) \to u_c^2/\kappa^2$ when $(x, y, u) \to (x_c, y_c, u_c)$.

7.1 Critical points

The critical points of the autonomous system (22)-(24) (with α replaced by $\alpha(u)$) are presented in table 3. In table 3 we define $\hat{\xi} \equiv 2\,\xi\,u_c^2 + 1$ and $\zeta \equiv \lambda + \beta > 0$. In table 4 we summarize the stability properties, and conditions for acceleration and existence for each point.

Point II.a corresponds to a scaling solution. The condition $0 \leq \Omega_\phi \leq 1$ is satisfied for $\beta < \sqrt{3/8}\,(2 - \gamma)$ if

$$\frac{3\,(2-\gamma)^2}{4\,\beta^2}\,\left(1 + \sqrt{1 - \frac{8\,\beta^2}{3\,(2-\gamma)^2}}\right) \leq \hat{\xi} \leq \frac{3\,(2-\gamma)^2}{2\,\beta^2}, \tag{48}$$

or

$$0 \leq \hat{\xi} \leq \frac{3\,(2-\gamma)^2}{4\,\beta^2}\,\left(1 - \sqrt{1 - \frac{8\,\beta^2}{3\,(2-\gamma)^2}}\right) < \frac{3\,(2-\gamma)^2}{2\,\beta^2}. \tag{49}$$

On the other hand, for $\beta \geq \sqrt{3/8}\,(2 - \gamma)$ it is required merely that

$$0 \leq \hat{\xi} \leq \frac{3\,(2-\gamma)^2}{2\,\beta^2}. \tag{50}$$

The equation of state for this solution is given by

$$\omega_\phi = \frac{2\,\beta^2\,(2-\gamma)\,\hat{\xi}}{\left(\hat{\xi}-1\right)\,\left(2\,\beta^2\,\left(\hat{\xi}-1\right) - 3\,(\gamma-4)\,\gamma + 4\,(\beta^2-3)\right) + 2\,\beta^2} + \gamma - 1. \tag{51}$$

For $\hat{\xi} \approx 1$ ($\xi \approx 0$), and $\beta > 0$, we have that $\omega_\phi \approx 1$. On the other when $\hat{\xi} \neq 1$ and $\beta = 0$, then $\omega_\phi \approx \gamma - 1 = \omega_m$.

The accelerated expansion occurs for

$$\hat{\xi} < -\frac{(2-\gamma)\,(3\,\gamma-2)}{2\,\beta^2} < 0. \tag{52}$$

Therefore the accelerated expansion no occurs in this solution since necessary $\hat{\xi}$ is positive to ensure $0 \leq \Omega_\phi \leq 1$.

Points II.b and II.c exists for $\hat{\xi} \geq 0$. Both points are scalar-field dominant solutions ($\Omega_\phi = 1$), but without accelerated expansion because $\omega_{eff} = 1$.

Point II.d is also a scaling solution. From the condition $0 \leq \Omega_\phi \leq 1$ we have the following constraint

$$\frac{3\,\gamma}{\lambda\,\zeta} \leq \hat{\xi} \leq \frac{3\,\gamma}{\lambda\,\zeta} + \frac{\zeta}{\lambda}. \tag{53}$$

The equation of state of field for this solution is given by

$$\omega_\phi = \frac{\beta\,\zeta\,\gamma}{\beta\,\lambda\,\left(\hat{\xi}-2\right) + \lambda^2\,\left(\hat{\xi}-1\right) - 3\,\gamma - \beta^2} + \gamma - 1, \tag{54}$$

and for $\gamma = 1$, $\hat{\xi} \approx 1$ and $\lambda << \beta$ ($\beta >> 1$) we have that $\omega_\phi \approx -1$ (De Sitter solution). The universe exhibits an accelerated expansion independently of $\hat{\xi}$ for

$$\lambda < \frac{2\,\beta}{3\,(\gamma-1)+1}. \tag{55}$$

In the case of non-relativistic dark matter ($\gamma = 1$) the accelerated expansion occurs for $\lambda < 2\,\beta$.

Point II.e is a scalar-field dominant solution ($\Omega_\phi = 1$) that exists for

$$0 \leq \hat{\xi} \leq \frac{6}{\lambda^2}, \tag{56}$$

and gives an accelerated expansion at late times for

$$\hat{\xi} < \frac{2}{\lambda^2}. \tag{57}$$

Given that $\hat{\xi} \geq 0$ is necessary for the existence of this solution, then $\omega_\phi = \omega_{eff} \geq -1$.

Table 3: Critical Points of the autonomous system (22)-(24) for dynamically changing $\alpha(u)$ such that $\alpha(u) \to \alpha(u_c) = 0$ and $u_c \geq 0$. We define $\hat{\xi} \equiv 2\,\xi\,u_c^2 + 1$ and $\zeta \equiv \lambda + \beta$.

Name	x_c	y_c	u_c	Ω_ϕ	ω_ϕ	ω_{eff}
II.a	$-\frac{\sqrt{6}\,\beta\,\hat{\xi}}{3\,(2-\gamma)}$	0	u_c	$\frac{2\,\beta^2\,\hat{\xi}^2}{3\,(\gamma-2)^2} + 1 - \hat{\xi}$	Eq. (51)	$\frac{2\,\beta^2\,\hat{\xi}}{3\,(2-\gamma)} + \gamma - 1$
II.b	$-\sqrt{\hat{\xi}}$	0	u_c	1	1	1
II.c	$\sqrt{\hat{\xi}}$	0	u_c	1	1	1
II.d	$\frac{\sqrt{6}\,\gamma}{2\,\zeta}$	$\frac{\sqrt{\beta\,\zeta\,\hat{\xi} + \frac{3}{2}\,(2-\gamma)\,\gamma}}{\zeta}$	u_c	$-\frac{\lambda\,\hat{\xi}}{\zeta} + \frac{3\,\gamma}{\zeta^2} + 1$	Eq. (54)	$\frac{\lambda\,(\gamma-1)-\beta}{\zeta}$
II.e	$\frac{\lambda\,\hat{\xi}}{\sqrt{6}}$	$\sqrt{\hat{\xi}\,\left(1 - \frac{\lambda^2\,\hat{\xi}}{6}\right)}$	u_c	1	$\frac{\lambda^2\,\hat{\xi}}{3} - 1$	$\frac{\lambda^2\,\hat{\xi}}{3} - 1$

Table 4: Stability properties, and conditions for acceleration and existence of the fixed points in table 3. As in table 3 we define $\hat{\xi} \equiv 2\,\xi\,u_c^2 + 1$ and $\zeta \equiv \lambda + \beta$.

Name	Stability	Acceleration	Existence
II.a	Saddle	No	Eqs. (48), (49) and (50)
II.b	Unstable node or saddle	No	$\hat{\xi} > 0$
II.c	Unstable node or saddle	No	$\hat{\xi} > 0$
II.d	Stable node or stable spiral or saddle	$\lambda < \frac{2\,\beta}{3\,(\gamma-1)+1}$	$\frac{3\,\gamma}{\lambda\,\zeta} \leq \hat{\xi} \leq \frac{3\,\gamma}{\lambda\,\zeta} + \frac{\zeta}{\lambda}$
II.e	Stable node or saddle	$\hat{\xi} < \frac{2}{\lambda^2}$	$0 \leq \hat{\xi} \leq \frac{6}{\lambda^2}$

7.2 Stability

For dynamically changing $\alpha(u)$ such that $\alpha(u) \to \alpha(u_c) = 0$, the components of the matrix of perturbation $\mathcal{M}$ are written as

$$\mathcal{M}_{11} = \frac{3\,\left(3\,(2-\gamma)\,x_c^2 - \gamma\,y_c^2\right)}{2\,\hat{\xi}} - \frac{3\,(2-\gamma)}{2} - \frac{\partial \hat{Q}}{\partial x}|_{x_c,y_c,u_c}, \tag{58}$$

$$\mathcal{M}_{12} = \sqrt{6}\,\lambda\,y_c - \frac{3\,\gamma\,x_c\,y_c}{\hat{\xi}} - \frac{\partial \hat{Q}}{\partial y}|_{x_c,y_c,u_c}, \tag{59}$$

$$\mathcal{M}_{13} = \frac{6\,\xi\,u_c\,x_c\,\left(\gamma\,y_c^2 - (2-\gamma)\,x_c^2\right)}{\hat{\xi}^2} + \frac{2\,\sqrt{6}\,\xi\,\eta_c\,u_c\,x_c^2}{\hat{\xi}} - \sqrt{6}\,\xi\,\eta_c\,u_c - \frac{\partial \hat{Q}}{\partial u}|_{x_c,y_c,u_c}, \tag{60}$$

$$\mathcal{M}_{21} = \frac{3\,(2-\gamma)\,x_c\,y_c}{\hat{\xi}} - \frac{\sqrt{6}\,\lambda\,y_c}{2}, \tag{61}$$

$$\mathcal{M}_{22} = \frac{3\,\left((2-\gamma)\,x_c^2 - 3\,\gamma\,y_c^2\right)}{2\,\hat{\xi}} - \frac{\sqrt{6}\,\lambda\,x_c - 3\,\gamma}{2}, \tag{62}$$

$$\mathcal{M}_{23} = \frac{6\,\xi\,u_c\,y_c\,\left(\gamma\,y_c^2 - (2-\gamma)\,x_c^2\right)}{\hat{\xi}^2} + \frac{2\,\sqrt{6}\,\xi\,\eta_c\,u_c\,x_c\,y_c}{\hat{\xi}}, \tag{63}$$

$$\mathcal{M}_{31} = 0, \qquad \mathcal{M}_{32} = 0, \qquad \mathcal{M}_{33} = \frac{\sqrt{6}\,\eta_c\,x_c}{2}. \tag{64}$$

Here we define $\eta_c \equiv \frac{d\eta(u)}{du}|_{u=u_c}$.

Using (32) in (58)-(60) we can calculate the components of $\mathcal{M}$ for each critical point and the corresponding eigenvalues.

- Point II.a:

$$\mu_1 = \frac{\beta\,\zeta\,\hat{\xi}}{2-\gamma} + \frac{3\,\gamma}{2}, \qquad \mu_2 = -\frac{\beta\,\eta_c\,\hat{\xi}}{2-\gamma}, \qquad \mu_3 = \frac{\beta^2\,\hat{\xi}}{2-\gamma} - \frac{3\,(2-\gamma)}{2}. \tag{65}$$

- Point II.b:

$$\mu_1 = -\frac{\sqrt{6}\,\eta_c\,\sqrt{\hat{\xi}}}{2}, \qquad \mu_2 = \frac{\sqrt{6}\,\left(\sqrt{6} + \lambda\,\sqrt{\hat{\xi}}\right)}{2}, \qquad \mu_3 = -\sqrt{6}\,\beta\,\sqrt{\hat{\xi}} + 3\,(2-\gamma)\,. \tag{66}$$

- Point II.c:

$$\mu_1 = \frac{\sqrt{6}\,\eta_c\,\sqrt{\hat{\xi}}}{2}, \qquad \mu_2 = \frac{\sqrt{6}\,\left(\sqrt{6} - \lambda\,\sqrt{\hat{\xi}}\right)}{2}, \qquad \mu_3 = \sqrt{6}\,\beta\,\sqrt{\hat{\xi}} + 3\,(2-\gamma)\,. \tag{67}$$

- Point II.d ($\gamma = 1$):

$$\mu_{1,2} = \frac{3\,(\lambda + 2\,\beta)}{4\,\zeta}\left(-1 \pm \sqrt{1 - \frac{8\,\left(2\,\beta\,\zeta\,\hat{\xi} + 3\right)\,\left(\zeta\,\lambda\,\hat{\xi} - 3\right)}{3\,(\lambda + 2\,\beta)^2\,\hat{\xi}}}\right), \qquad \mu_3 = \frac{3\,\eta_c}{2\,\zeta}. \tag{68}$$

- Point II.e:

$$\mu_1 = \lambda\,\zeta\,\hat{\xi} - 3\,\gamma, \qquad \mu_2 = \frac{\lambda^2\,\hat{\xi} - 6}{2}, \qquad \mu_3 = \frac{\eta_c\,\lambda\,\hat{\xi}}{2}. \tag{69}$$

Since for the point II.a we have $0 \le \hat{\xi} \le 3\,(2-\gamma)^2/2\,\beta^2$ (Eqs. (48), (49) and (50)), then it is always a saddle point. Point II.b can be unstable node or saddle point. For $0 < \hat{\xi} < 3\,(2-\gamma)^2/2\,\beta^2$ and $\eta_c < 0$ it is unstable node. For $\eta_c > 0$ or $\hat{\xi} > 3\,(2-\gamma)^2/2\,\beta^2$ it is saddle point. Point II.c is a unstable node if $0 < \hat{\xi} < 6/\lambda^2$ and $\eta_c > 0$, whereas for $\eta_c < 0$ or $\hat{\xi} > 6/\lambda^2$ it is a saddle point.

To study the properties of stability of the point II.d we shall consider the case where the background fluid is non-relativistic dark matter ($\gamma = 1$). So, we restrict ourselves to positive $\hat{\xi}$ as in constraint (53) with $\gamma = 1$, and $\lambda < 2\,\beta$ or equivalently $\zeta < 3\,\beta$.

The eigenvalues μ_1 and μ_2 are real and negatives if

$$\frac{3}{\zeta\,\lambda} < \hat{\xi} \leq \Delta + \frac{3}{\zeta\,\lambda}, \tag{70}$$

where

$$\Delta = \frac{3\left(\sqrt{\delta^2 + 64\,\beta\,\zeta\,(\lambda + 2\,\beta)^2} + \delta\right)}{32\,\beta\,\zeta^2\,\lambda} > 0, \tag{71}$$

with $\delta = \left(-7\,\zeta^2 - 6\,\beta\,\zeta + \beta^2\right)$. Then, for $\eta_c < 0$ and $\hat{\xi}$ satisfying (70) the point II.d is a stable node. For $\eta_c > 0$ and $\hat{\xi}$ as in (70) it is saddle point. On the other hand for

$$\Delta + \frac{3}{\zeta\,\lambda} < \hat{\xi} \leq \frac{3}{\lambda\,\zeta} + \frac{\zeta}{\lambda}, \tag{72}$$

μ_1 and μ_2 are complex with real part negative, and the determinant of the matrix $\mathcal{M}$ that is given by

$$\det(\mathcal{M}) = \frac{9\,\eta_c\left(2\,\beta\,\zeta^3\,\lambda\,\hat{\xi}^2 + 3\,\zeta^2\,(\lambda - 2\,\beta)\,\hat{\xi} - 9\right)}{4\,\zeta^4\,\hat{\xi}}, \tag{73}$$

is negative if in addition $\eta_c < 0$ (also in this case $\mu_3 < 0$). Therefore for $\hat{\xi}$ as in (72) and $\eta_c < 0$, point II.d becomes stable spiral. The constraint (70) is consistent with the physical condition (53) (it can be stronger or weaker). On the other hand, the constraint (72) is also consistent with (53) provided that $\Delta < \zeta/\lambda$. In any case (stable node or stable spiral), point II.d.2 is a scaling attractor with accelerated expansion ($\lambda < 2\,\beta$).

Point II.e is a stable node if

$$0 < \hat{\xi} < \frac{3\,\gamma}{\lambda\,\zeta} < \frac{6}{\lambda^2}, \tag{74}$$

and $\eta_c < 0$. Since $1 \leq \gamma < 2$ then $\frac{3\,\gamma}{\lambda\,\zeta} < \frac{6}{\lambda^2}$. On other hand, it is a saddle point for

$$\frac{3\,\gamma}{\lambda\,\zeta} < \hat{\xi} < \frac{6}{\lambda^2}. \tag{75}$$

or $\eta_c > 0$. The point II.e is also an interesting solution, it is a scalar-field dominant solution ($\Omega_\phi = 1$) and can be attractor (Eq. (74)) with accelerated expansion (Eq. (57)) and such that $\omega_\phi = \omega_{eff} \geq -1$.

So, for dynamically changing α such that $\alpha(u) \to \alpha(u_c) = 0$, the fixed points II.d and II.e can attract the universe at late times and are equally viable solutions. But since we are interested in the scaling attractors, now we are going to concentrate on the solution II.d. It is a scaling solution that has accelerated expansion and at the same time is an attractor (stable node or stable spiral). For $\gamma = 1$, the accelerated expansion occurs, independently of $\hat{\xi}$, for $\lambda < 2\,\beta$ (Eq. (55)), and the physical condition $0 \leq \Omega_\phi \leq 1$ is ensures for $\hat{\xi} > 0$ in accordance with (53). Also, for $\hat{\xi}$ in accordance with (70) and $\eta_c < 0$ point II.d is a stable node, or for $\hat{\xi}$ as in (72) and $\eta_c < 0$ it is a stable spiral. When point II.d is a stable node, the constraint (70) can be stronger or weaker that the physical condition (53). Numerically we find that exist



Figure 1: Evolution of Ω_m (dashed), Ω_ϕ (dotdashed), ω_ϕ (dotted), ω_{eff} (solid) and $\alpha(u)$ (red line) with $\gamma = 1$, $\lambda = 2.4$ and $\xi \approx 3.5 \times 10^{-5}$, in the case where the coupling to dark matter β changes rapidly from $\beta_1 = 0$ to $\beta = 4.1$ ($b = 0.035$ and $u_1 = 0.21$). We choose initial conditions $x_i = 5 \times 10^{-5}$, $y_i = 2.6 \times 10^{-6}$ and $u_i = 7 \times 10^{-4}$ and by way of example we consider the function $\alpha(u) = u_c - u$ with $u_c = 1$ and $\eta_c = -1$. The universe exits from the matter-dominated solution I.a with constant α, $\omega_\phi = -1$, $\omega_{eff} = 0$ and approaches the scaling attractor II.d for dynamically changing $\alpha(u)$ with $\Omega_\phi \approx 0.70$, $\omega_\phi \approx -0.90$ and $\omega_{eff} \approx -0.63$. We note that ω_ϕ presents the phantom-divide crossing during cosmological evolution.

$\lambda_*(\beta) < 2\,\beta$ such that for $\lambda < \lambda_*(\beta)$ then $\Delta > \zeta/\lambda$ and the constraint (70) is weaker that (53). For example, for $\beta = 0.6$ then $\lambda_*(\beta) \approx 0.82$, and if $\lambda = 0.2$ the constraint (70) implies $18.75 < \hat{\xi} < 22.96$. Moreover, the physical condition (53) implies $18.75 < \hat{\xi} < 22.75$. Also, in this example, for $\hat{\xi} \approx 19.95$ then $\omega_\phi \approx -1.07$ (Eq. (54)) and $\Omega_\phi \approx 0.70$ (with accelerated expansion since $\lambda < 2\,\beta$). However, this is not a suitable range for $\hat{\xi}$, since from solar system experiments (see for instance [6]), necessary $\hat{\xi} \approx 1$ (or $\xi \approx 0$). Finally, when point II.d is a stable spiral, that is, $\hat{\xi}$ in accordance with (72) and $\eta_c < 0$, the physical condition (53) is ensures whenever $\Delta < \zeta/\lambda$. For example, for $\lambda = 2.4$ and $\beta = 4.1$ we have that (72) implies $0.23 < \hat{\xi} < 2.90$, whereas the constraint (53) implies $0.19 < \hat{\xi} < 2.90$. In Fig. 1 we show the case when the system approaches the fixed point II.d in the above example for $\lambda = 2.4$, $\beta = 4.1$ and for instance $\hat{\xi} \approx 1.00064$ ($\xi \approx 3.5 \times 10^{-5}$ and $u_c = 1$). In this figure, by way of example, we consider the function $\alpha(u) = u_c - u$ with $u_c = 1$ and $\eta_c = -1$. Also, following [1, 31], we consider a non-linear coupling

$$\beta(u) = \frac{(\beta - \beta_1)\,\tanh\left(\frac{u-u_1}{b}\right) + \beta + \beta_1}{2}, \tag{76}$$

that changes between a small β_1 to a large β in order to that initially the system enters in the matter-dominated solution I.a (constant α) with $\beta_1 = 0$ and eventually approaches the scaling attractor II.d with $\omega_\phi \approx -0.90$, $\Omega_\phi \approx 0.70$ and accelerated expansion ($\lambda < 2\,\beta$).

8 Conclusions

It has been recently proposed [15, 16] a non-minimal coupling ϕ^2 between quintessence and gravity in the framework of teleparallel gravity, motivated by a similar construction in the context of GR. The theory has been called “teleparallel dark energy”, and the cosmological dynamics of the model was studied later in [17, 18], but no scaling attractor was found.

In this paper we generalize the function of non-minimal coupling $\phi^2 \to f(\phi)$ and we define $\alpha \equiv f_{,\phi}/\sqrt{f}$ such that α is constant for $f(\phi) \propto \phi^2$, but it can also be a dynamically changing quantity $\alpha(\phi)$. For constant $\alpha > 0$ and ξ as in (47) ($\beta = 0$), the universe enters the matter-dominated solution I.a ($\Omega_m = 1$) with $\omega_\phi = -1$ and $\omega_{eff} = 0$, but since this solution is unstable, the universe will eventually fall into the attractor I.d (dark-energy-dominated de Sitter solution) with $\Omega_\phi = 1$ and $\omega_\phi = \omega_{eff} = -1$. Also, for constant α no scaling attractor was found, even when we allow a possible interaction between teleparallel dark energy and dark matter.

On other hand, for dynamically changing $\alpha(\phi)$, and when it can be expressed in terms of $u \equiv \kappa\sqrt{f(\phi)}$ such that $\alpha(u) \to \alpha(u_c) = 0$ (we notice that (x_c, y_c, u_c) is a fixed point of the system), we show that interesting solutions exist (see table 3), in particular scaling attractors with accelerated expansion. These are non-minimal generalizations of the fixed points presented in [1, 30, 32] for coupled dark energy in GR. The final attractor is either the scalar-field dominated solution II.e or the scaling attractor II.d, both with accelerated expansion. The fixed point II.d is a scaling attractor that can be a stable node or stable spiral when the background fluid is non-relativistic dark matter ($\gamma = 1$). When point II.d is a stable spiral, that is, $\hat{\xi} \equiv 2\,\xi\,u_c^2 + 1$ as in constraint (72), and

$$\eta_c \equiv \frac{d\alpha(u)}{du}|_{u=u_c} < 0,$$

it is possible to find a range for $\hat{\xi}$ in accordance with solar system experiments ($\hat{\xi} \approx 1$) [6], and compatible with cosmological observations. For example, in Fig. 1 the system approaches point II.d with $\Omega_\phi \approx 0.70$, $\Omega_m \approx 0.30$, $\omega_\phi \approx -0.90$ and accelerated expansion, in agreement with observations. Also, we note that ω_ϕ presents the phantom-divide crossing during cosmological evolution, as has been stressed in Refs. [15, 16, 18] for teleparallel dark energy. Since the scaling solution II.d attracts the universe at late times regardless of the initial conditions with Ω_ϕ comparable to Ω_m and accelerated expansion, the cosmological coincidence problem could be alleviated without fine-tunings [1, 4].

Acknowledgments

The author would like to thank J. G. Pereira and E.N. Saridakis for useful discussions and suggestions. He would like to thank also CAPES for financial support.

References

[1] E.J. Copeland, M. Sami, S. Tsujikawa, Int. J. Mod. Phys. D 15 (2006) 1753, hep-th/0603057;

J. Frieman, M. Turner, D. Huterer, Ann. Rev. Astron. Astrophys. 46 (2008) 385, arXiv:0803.0982;

S. Tsujikawa, arXiv:1004.1493;

M. Li, X.D. Li, S. Wang, Y. Wang, Commun. Theor. Phys. 56 (2011) 525, arXiv:1103.5870.

[2] A. De Felice, S. Tsujikawa, Living Rev. Rel. 13 (2010) 3, arXiv:1002.4928;

T. Clifton, P.G. Ferreira, A. Padilla, C. Skordis, arXiv:1106.2476.

[3] T.P. Sotiriou, V. Faraoni, Rev. Mod. Phys. 82 (2010) 451, arXiv:0805.1726;

S. Tsujikawa, Lect. Notes Phys. 800 (2010) 99, arXiv:1101.0191;

S. Capozziello, M. De Laurentis, V. Faraoni, arXiv:0909.4672;

R. Durrer, R. Maartens, arXiv:0811.4132.

[4] L. Amendola, S. Tsujikawa, *Dark energy, theory and observations* (Cambridge Univ. Press, Cambridge, 2010).

[5] F. Perrotta, C. Baccigalupi and S. Matarrese, Phys. Rev. D 61 (2000) 023507, astro-ph/9906066.

[6] V. Sahni, S. Habib, Phys. Rev. Lett. 81 (1998) 1766, hep-ph/9808204;

T. Chiba, Phys. Rev. D60 (1999) 083508, gr-qc/9903094;

J.P. Uzan, Phys. Rev. D 59 (1999) 123510, gr-qc/9903004;

N. Bartolo, M. Pietroni, Phys. Rev. D 61 (2000) 023518, hep-ph/9908521;

V. Faraoni, Phys. Rev. D 62 (2000) 023504, gr-qc/0002091;

E. Elizalde, S. Nojiri, S.D. Odintsov, Phys. Rev. D 70 (2004) 043539, hep-th/0405034.

[7] O. Hrycyna, M. Szydlowski, JCAP 0904 (2009) 026, arXiv:0812.5096;

O. Hrycyna, M. Szydlowski, Phys. Rev. D 76 (2007) 123510, arXiv:0707.4471.

[8] R. C. de Souza, G. M. Kremer, Class. Quant. Grav. 26 (2009) 135008, arXiv:0809.2331.

[9] A.A. Sen, N. Chandrachani Devi, Gen. Rel. Grav. 42 (2010) 821, arXiv:0809.2852.

[10] K. Hayashi, T. Shirafuji, Phys. Rev. D 19 (1979) 3524.

[11] J.W. Maluf, J. Math. Phys. 35 (1994) 335;

H.I. Arcos, J.G. Pereira, Int. J. Mod. Phys. D 13 (2004) 2193, gr-qc/0501017.

[12] R. Aldrovandi, J. G. Pereira, *Teleparallel Gravity: An Introduction* (Springer, Dordrecht, 2013);

J. G. Pereira, *Lorentz connections and gravitation*, AIP Conf. Proc. 1483 (2012) 239-259, arXiv:1210.0379;

J. G. Pereira, *Teleparallelism: a new insight into gravitation*, in *Springer Handbook of Spacetime*, ed. by A. Ashtekar and V. Petkov (Springer, Dordrecht, 2013) arXiv:1302.6983.

[13] G.R. Bengochea, R. Ferraro, Phys. Rev. D 79 (2009) 124019, arXiv:0812.1205.

[14] E.V. Linder, Phys. Rev. D 81 (2010) 127301, arXiv:1005.3039.

[15] Chao-Qiang Geng, Chung-Chi Lee, E.N. Saridakis, Yi-Peng Wu, Phys. Lett. B 704 (2011) 384, arXiv:1109.1092.

[16] Chao-Qiang Geng, Chung-Chi Lee, E.N. Saridakis, JCAP 1201 (2012) 002, arXiv:1110.0913.

[17] Hao Wei, Phys. Lett. B 712 (2012) 430, arXiv:1109.6107.

[18] Chen Xu, E.N. Saridakis, G. Leon, JCAP 1207 (2012) 005, arXiv:1202.3781.

[19] F.S. Accetta, D.J. Zoller, M.S. Turner, Phys. Rev. D 31 (1985) 3046.

[20] L. Amendola, M. Litterio, F. Occhionero, Int. J. Mod. Phys. A 5 (1990) 3861.

[21] T. Futamase, K. Maeda, Phys. Rev. D 39 (1989) 399.

[22] U. Kasper, Nuovo Cim. B 103 (1989) 291.

[23] P.J. Steinhardt, F.S. Accetta, Phys. Rev. Lett. 64 (1990) 2740.

[24] L. Amendola, Phys. Rev. D 60 (1999) 043501, astro-ph/9904120.

[25] L. Amendola, Phys. Lett. B 301 (1993) 175, gr-qc/9302010;
S.F. Daniel, R.R. Caldwell, Class. Quant. Grav. 24 (2007) 5573, arXiv:0709.0009;
E.N. Saridakis, S.V. Sushkov, Phys. Rev. D 81 (2010) 083510, arXiv:1002.3478;
C. Germani, A. Kehagias, Phys. Rev. Lett. 105 (2010) 011302, arXiv:1003.2635.

[26] R.J. Scherrer, A. Sen, Phys. Rev. D 77 (2008) 083515, arXiv:0712.3450.

[27] R.J. Scherrer, A. Sen, Phys. Rev. D 78 (2008) 067303, arXiv:0808.1880.

[28] M. Setare, E.N. Saridakis, Phys. Rev. D 79 (2009) 043005, arXiv:0810.4775.

[29] Shuang-Yong Zhou, Phys. Lett. B 660 (2008) 7, arXiv:0705.1577;
Wei Fang, Ying Li, Kai Zhang, Hui-Qing Lu, Class. Quant. Grav. 26 (2009) 155005, arXiv:0810.4193.

[30] B. Gumjudpai, T. Naskar, M. Sami, S. Tsujikawa, JCAP 0506 (2005) 007, hep-th/0502191.

[31] L. Amendola, D. Tocchini-Valentini, Phys. Rev. D 64 (2001) 043509, astro-ph/0011243.

[32] L. Amendola, Phys. Rev. D 62 (2000) 043511, astro-ph/9908023.